## Supplemental Labeling



DowElanco Quad IV, 9002 Purdue Road

P.O. Box 681428

Indianapolis, Indiana 46268-1189 USA

TREFLAM\* TR-10 (E.P.A. Reg. No. 62719-131) (For Distribution and Use Only in the State of Idaho) 24(c) Special Local Need Registration (SLN ID-91 0001)

Treflan TR-#0 For Dodder Control in Established Alfalfa Grown for Seed

This labeling must be in the possession of the user at the time of application.

Under this Special Local Need Registration TREFLAN TR-10 may be surface applied in established alfalfa grown for seed only, at 30 to 40 pounds per acre. Treated alfalfa seed fields must not be used for human or animal consumption and use of alfalfa forage, hay or seed byproducts for food or feed is prohibited (see "Use Restrictions" below).

Application Directions Apply at a broadcast rate of 30 to 40 pounds per acre in late winter or spring. Application and activation must occur before dodder germination. For best results TREFLAN TR-10 should be activated within 3 days after application. When surface applied, TREFLAN TR-10 may be activated by rainfall, overhead sprinkler irrigation, flood irrigation or furrow irrigation. If activated using furrow irrigation, the surface of beds between furrows should be thoroughly wetted.

Use Restrictions: The following use restrictions apply to alfalfa fields treated with TREFLAN TR-10 as described above.

- 1. For use on fields in the production of alfalfa seed. Not for use on fields producing alfalfa for livestock feed. No portion of the treated field, including seed, seed screenings, hay, forage or stubble, may be used for human or animal consumption.
- 2. All seed screenings must be disposed of in such a way that they cannot be recovered and used for food or feed.
- 3. Do not use alfalfa seed for sprouting. Processed seed must be labeled "Not for human or animal consumption" at the processing plant.
- 4 Use of this product according to this labeling is considered a non-food use.

Use precautions:

- Where alfalfa is to be sotated to another crop in the season following application of TREFLAN TR-10, plant only those crops to which TREFLAN may be applied as a preplant incorporated reatment or crop injury may occur. Read and carefully follow all rotational crop restrictions on the TREFLAN TR-10 label. The potential for rotational crop injury may be reduced by moldboard plowing to a depth of 12 inches prior to planting.
- Do not apply to seedling alfalfa (up to 6 true leaf stage).
- Refer to the TREFLAN TR-10 label for additional directions, precautions and limitations prior to use.

\*Trademark of DowElanch 00/00/00 T2EAGXXX Approved

This is an Initial DowElanco Printing.

The second secon